

# Paper 2

*by* Rachmaniyah 2

---

**Submission date:** 06-Nov-2019 08:22AM (UTC+0700)

**Submission ID:** 1207928927

**File name:** ractas\_Antioxidant\_on\_Cigarette\_Smoke\_Exposure\_in\_Mice\_lungs.pdf (104.5K)

**Word count:** 2199

**Character count:** 12394



Scientia Research Library

ISSN 2348-0416

USA CODEN: JASRHB

Journal of Applied Science And Research, 2018, 6 (5):11-15

(<http://www.scientiaresearchlibrary.com/archive.php>)

## The Potency of Red Guava(*Psidium Guajava L.*)Extractas Antioxidant on Cigarette Smoke Exposure in Mice lungs

Imam Thohari<sup>1</sup>, Rachmaniyah<sup>1</sup>, Umi Rahayu<sup>1</sup>

<sup>1</sup>Study Program of Environmental Health, Polytechnic of Health, Surabaya 60115 Indonesia

### ABSTRACT

Cigarette smoke can cause to the formation of an oxidative stress, resulting in increase in concentration of free radicals and decrease antioxidant. *Psidium guajava* L extract has shown to possess powerful antioxidant properties. In the present study, we investigated the impact of *Psidium guajava* L extract in against cigarette smoke-induced lung damage in mice. The 28 male mice were divided into: negative control (mice were exposed with fresh air daily 1 h twice a day for 30 days); positive control (mice were exposed with cigarette smoke 1 h twice a day for 30 days), and the treatment group (mice were given the *Psidium Guajava* L extract 0.3 ppm, 0.70 ppm orally once in a day for 34 days and on 4<sup>th</sup> day, were exposed with cigarette smoke 1 h twice a day for 30 days. On day 34, the mice were sacrificed, and lung tissues were collected to histological evaluations. The cigarette smoke significantly induced necrosis in lung cell. However, treatment with the *Psidium Guajava* L extract significantly improved lung cell damage. From the results, it is concluded that the *Psidium Guajava* L extract is a potent antioxidant against cigarette smoke-induced lung damage in mice.

**Keywords :** *Psidium Guajava L.*, Antioxidant, Lung, Cigarette smoke

### INTRODUCTION

Cigarette smoke (CS), with its over 4700 different substances (Genbacev-Krtolica, 2005), most of which are described as toxic, mutagenic and carcinogenic (Czekaj et al., 2002), is involved in the development of various chronic pulmonary, cerebro vascular and cardio vascular diseases, cancers and several others (Pryor, 1997). Harmful influences of cigarette smoke can be mainly associated with its ingredients like nicotine, cadmium, benzopyrene, oxidants, acetone, mercury and inducers of reactive oxygen species (ROS) like nitric oxide (NO), nitrogen dioxide (NO<sub>2</sub>), peroxynitrite and nitroamines. The number of ROS, especially O<sub>2</sub><sup>-</sup>, hydroxyl radical (OH) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) increases after CS exposure in cells (Rahman, 2005). Reactive oxygen species have a considerable role in smoking related diseases by initiating lipid peroxidation (LPO) and protein oxidation (Reznick et al., 1992) as well as causing DNA strand breaking (Kodama et al., 1997). One puff of cigarette smoke includes 1015 oxidant radicals (Mc Cusker and Hoidal, 1990).

Reactive oxygen species are suppressed or scavenged by cellular antioxidant defense systems

consisting of enzymatic mechanisms such as super oxide <sup>13</sup> dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx) and glutathione reductase (GR) as well as non-enzymatic mechanisms such as <sup>9</sup> reduced glutathione (GSH), vitamins A, C and E. Preventive role of antioxidants was reported in modifying the major diseases related to Cigarette smoke (Anbarasi et al., 2005). When ROS increased excessively, antioxidant deficiency occurs. This inadequacy may be enhanced by usually insufficient intake of both supplementary and dietary antioxidants (Zondervan et al., 1996). One of the materials being used as antioxidant source to eliminate free radicals in CS are plant extracts (Chang et al., 2010).

Antioxidant was mostly in plant, and one of natural antioxidant sources was red guava (*Psidium guajava* L) which was very nutritious because it contained ascorbic acid (50-3000mg)/100g fresh weight), three until six times higher rather than orange (Agustina, 2009). Red guava (*Psidium Guajava* L) was a fruit with vitamin C and ascorbic acid even flavonoid compound which had antioxidant that could slow down aging process and damage of the membrane. The objective of the present study was to investigate the antioxidant activity of the *Psidium Guajava* L extract on cigarette smoke-induced lung cell damage in mice.

## MATERIAL AND METHODS

### Preparation of ethanol extract of *Psidium Guajava* L

<sup>3</sup> Plant material and extract preparation <sup>3</sup> *Psidium Guajava* L were collected from Surabaya, Indonesia. *Psidium Guajava* L materials were cleaned with running tap water and chopped into pieces. They were dried under shade at ambient temperature for 5 days, and the air dried *Psidium Guajava* L was then ground to powder for extraction. The powdered *Psidium Guajava* L (1 kg) was <sup>7</sup>acerated with ethanol 96% (5 L) for 1 week at 37°C. The supernatant was then collected and filtered through Whatman No. 1 filter paper in a Buchner funnel under vacuum. The filtrate was concentrated by evaporation with a vacuum rotary evaporator at 45°C. Further more, the extract was freeze dried to get ethanol-free powder.

### Experimental animal

Male mice weighing approximately 24-30 g (2.5-3 months) <sup>19</sup> were obtained from Airlangga University, Surabaya, Indonesia for experimental purpose. They were housed in plastic cages in an air-conditioned room with a <sup>18</sup> temperature maintained at  $26 \pm 2$  °C and 12 h alternates light and dark cycles. The mice were given ad libitum with tap water and fed with standard commercial rat chow.

### Experimental design


The sample used <sup>21</sup> male mice were divided into 4 groups: negative control (mice were exposed with fresh air daily 1 h twice a day for 30 day); positive control (Mice were exposed with cigarette smoke 1 h twice a day for 30 days), and the treatment group (mice were given the *Psidium Guajava* L extract 0.35 ppm, 0. <sup>10</sup> ppm orally once in a day for 34 days and on 4<sup>th</sup> day, were exposed with cigarette smoke 1 h twice a day for 30 days. On day 34, the mice were sacrificed <sup>20</sup> and lung tissues were collected to histological evaluations. The tissue of lung was fixed in a 10% ne <sup>20</sup> al buffered formalin solution, embedded in paraffin and used for histopathological examination with hematoxylin and eosin (H&E) stain.

## RESULT AND DISCUSSION

Histological observations on the negative control showed that in lung cell is observable and they appear normal architecture of the lung cell. In the positive control, mice were administered with cigarette smoke showed lung cell damage (necrosis). In the mice, treated with *Psidium Guajava* L



extract, the number and morphological integrity of lung cells are being preserved. Observations indicate that the damage effects cigarette smoke was reduced by *Psidium Guajava* L extract [Figure 1].



**Figure 1:** Histological study of treatment with *Psidium Guajava* L extract on cigarette smoke-caused lung damage. Normal morphology of lung sections in the negative control group (A). The cigarette smoke treated group showed necrosis (indicated by black arrows) (B). Mice treated with *Psidium Guajava* L extract 0.35 ppm and 0.75 ppm showed necrotic changes (C and D). H&E  $\times 400$

## DISCUSSION

Cigarette smoke includes many oxidants and pro-oxidants that generate free radicals and increase the oxidative stress in lung tissues (Park et al., 1998). Thus, cigarette smoke is a constant oxidant stress on the respiratory tract. A cell could protect itself against oxidative stress by the use of endogenous and exogenous antioxidants. Depending on this data, smokers are usually directed to use antioxidants (Smith and Hodges, 1987). Current study was realized to determine whether *Psidium Guajava* L extract would be an effective antioxidant against smoking-mediated oxidative damage.

Cigarette smoke exposure was a trigger of necrosis based on that was influenced by the duration of smoking, the number of cigarettes smoked per day, the age and depth of smoking. Cigarette smoke could lead the entry of free radicals in the body and cause necrosis on epithelium. The process of necrosis began with the weakening of blood vessel wall, inflammation response, the formation of plaque that was foam cells and the formation of blood clots. Cigarette smoke exposure for a month aimed to provide allergen exposure directly to the main target cause the occurrence of bronchitis. Normal respiratory tract wall was layered by ciliated pseudo stratified. The condition of bronchitis caused cell changes and structural abnormalities due to inflammation. Through histopathologic examination on bronchus, it was obtained damage of epithelial bronchus with cilia abnormalities and smooth muscle hypertrophy.

Necrosis could be recognized due to the changes either macroscopically or microscopically. Necrosis caused nucleus cell changes that consisted of 3 processes of picnosis<sub>1</sub>, which was core contraction that was occurred due to homogenizing cytoplasm and the increase of eosinophil, and then, DNA condensed into a dense mass. After that, it was occurred cariorexis<sub>2</sub>, which was a pignotic fragmented nit state (divided into fragments), then, it was occurred cariolisis<sub>3</sub>, which was the fading of basophil chromatin due to DNA se activity (Lestari and Mulyono 2011).

Psidium Guajava L. as an antioxidant contained various substances that had a function as an inhibitor of various diseases, including flavonoids, tannins, oil and also Saponin. Flavonoid was an alcohol in the form of Phenol. Meanwhile, polyphenol compounds of red guava were quercetin, guajavarin, gallic acid, leukocyanidin 0.1% hexahydroxydiphenyl ester in the form of glycoside 0.1%, elagat acid (Agustina, 2009).

Flavonoid could prevent LDL oxidation 20 times stronger rather than vitamin C. Flavonoids were proven to have very strong biological effects as anti-oxidants that stimulated the production of nitrite that could widen blood vessels. Free radicals in the body affected the work of endogenous antioxidants. Continuous cigarette smoke exposure could increase free radicals in the body that would trigger an imbalance between free radicals and endogenous antioxidants that caused oxidative stress. Flavonoid in red guava extract (Psidium guajava L) could increase endogenous antioxidant activity (SOD, Catalase, GPx), thus, it could stabilize free radical bonds in the body, and the IgI decreased. The improvement process of epithelial was occurred by the decrease of epithelial damage when the free electrons from cigarette smoke exposure were tied to flavonoid antioxidant of red guava extract (Psidium guajava L) in the form of OH cluster. Red guava extract (Psidium Guajava L) would trigger SOD, Catalase and GPx activity in the body in order to stabilize free radicals. Nevertheless, the results showed administration of Psidium Guajava L extract could improve lung tissue damage (inflammation, oedema, necrosis) that was caused by cigarette smoke exposure.

## CONCLUSION

The results of the present study indicate that cigarette smoke-caused lung damage (necrosis) might be related to oxidative stress. The administration of Psidium Guajava L extract lessened the effects of cigarette smoke-induced lung tissue necrosis possibly by inhibiting free radical-mediated process. Further investigation of these promising protective effects of Psidium Guajava L extract against cigarette-induced lung cell damage may have a considerable impact on developing clinically feasible strategies to treat patients with cigarette smoke-induced lung damage.

## REFERENCES

- [1] Agustina R. (2009). The Effect of Giving Ethanol Extract in 70% from the Inside of Red Guava (*Psidium Guajava L.*) against cholesterol content in the serum of Wistar Male White Mice's Blood, *Rattus Norvegicus*. Thesis. Muhammadiyah University of Surakarta.
- [2] Anbarasi K, Sabitha KE, and Devi CSS (2005) Lactate dehydrogenase isoenzyme patterns upon chronic exposure to cigarette smoke: Protective effect of Bacoside A. *Environmental Toxicology and Pharmacology* 20: 345–350.
- [3] Chang GW, Dai Y, Dong LL, and Kuo YM (2010) Ginkgo biloba leaf extract action in scavenging free radicals and reducing mutagenicity and toxicity of cigarette smoke in vivo. *Journal of Environmental Science and Health* 45:498–505.
- [4] Czekaj P, Palasz A, Lebda-Wyborny T, et al. (2002) Morphological changes in lungs, placenta, liver and kidneys of pregnant rats exposed to cigarette smoke. *International Archives of Occupational and Environmental Health* 75: 27–35.
- [5] Genbacev-Krtolica O (2005) Highlight for phenols, quinolines, indoles, benzene and 2 cyclopenten-1-ones are oxidant oxicans in cigarette smoke, by Prue Talbot, Karen Riveles, and Ryan Rosa: List of tobacco-smoke constituents that are harmful for reproduction grows-passive smokers may be at risk. *Toxicological Sciences* 86: 4–5.

- [6] <sup>5</sup> Kodama M, Kaneko M, Aida M, Inoue F, Nakayama T, and Akimoto H (1997) Free radical chemistry of cigarette smoke and its implication in human cancer. *Anticancer Research* 17: 433–437.
- [7] Lestari SP, Mulyono A (2011). Analysis of Kidney for identifying either pyknosis cell or necrosis cell. *Journal of Neutrino* (1):48-66
- [8] <sup>6</sup> McCusker K and Hoidal J (1990) Selective increase of antioxidant enzyme activity in the alveolar macrophages from cigarette smokers and smoke-exposed hamsters. *American Review of Respiratory Disease* 141: 678–682.
- [9] <sup>8</sup> Park EM, Park YM, and Gwak YS (1998) Oxidative damage in tissues of rats exposed to cigarette smoke. *Free Radical Biology and Medicine* 25: 79–86.
- [10] <sup>14</sup> Pryor WA (1997) Cigarette smoke radicals and the role of free radicals in chemical carcinogenesis. *Environmental Health Perspectives* 105: 875–882.
- [11] <sup>11</sup> Rahman I (2005) The role of oxidative stress in the pathogenesis of COPD: Implications for therapy. *Treatments in Respiratory Medicine* 4: 175–200.
- [12] <sup>10</sup> Reznick AZ, Cross CE, Hu ML, et al. (1992) Modification of plasma proteins by cigarette smoke as measured by protein carbonyl formation. *Biochemical Journal* 286:607-611.
- [13] Smith JL and Hodges RE (1987) Serum level of vitamin C in relation to dietary and supplemental intake of vitamin C in smokers and nonsmokers. *Annals of the New York Academy of Sciences* 498: 443–447.
- [14] Zondervan KT, Ocke MC, Smit HA, and Seidell J (1996) Do dietary and supplementary intakes of antioxidants differ with smoking status? *International Journal of Epidemiology* 25: 70–79.



## Paper 2

### ORIGINALITY REPORT

23%

SIMILARITY INDEX

14%

INTERNET SOURCES

20%

PUBLICATIONS

10%

STUDENT PAPERS

### PRIMARY SOURCES

1

T. H. Thatcher. "Role of CXCR2 in cigarette smoke-induced lung inflammation", AJP Lung Cellular and Molecular Physiology, 03/11/2005

Publication

3%

2

[insights.ovid.com](https://www.insights.ovid.com)

Internet Source

2%

3

[www.florajournal.com](http://www.florajournal.com)

Internet Source

2%

4

Pekmez, H., M. Ogeturk, H. Ozyurt, M. F. Sonmez, N. Colakoglu, and I. Kus. "Ameliorative effect of caffeic acid phenethyl ester on histopathological and biochemical changes induced by cigarette smoke in rat kidney", Toxicology and Industrial Health, 2010.

Publication

1%

5

[www.fasebj.org](http://www.fasebj.org)

Internet Source

1%

6

Submitted to The University of Manchester

Student Paper

1%

7	<a href="http://www.abpv.vet.br">www.abpv.vet.br</a> Internet Source	1 %
8	Submitted to 89986 Student Paper	1 %
9	G. R. Thome. "Vitamin E Decreased the Activity of Acetylcholinesterase and Level of Lipid Peroxidation in Brain of Rats Exposed to Aged and Diluted Sidestream Smoke", Nicotine & Tobacco Research, 12/01/2011 Publication	1 %
10	Frayn. "References", Cardiovascular Disease, 04/04/2005 Publication	1 %
11	<a href="http://metabiolab.com">metabiolab.com</a> Internet Source	1 %
12	Submitted to Wartburg College Student Paper	1 %
13	<a href="http://www.scielo.br">www.scielo.br</a> Internet Source	1 %
14	<a href="http://who.int">who.int</a> Internet Source	1 %
15	Talib Hussain, Ramesh K Gupta, K Sweety, Bavani Eswaran, M Vijayakumar, Chandana Venkateswara Rao. "Nephroprotective activity of Solanum xanthocarpum fruit extract against	1 %



gentamicin–induced nephrotoxicity and renal dysfunction in experimental rodents", Asian Pacific Journal of Tropical Medicine, 2012

Publication

16

Submitted to The New Art College

Student Paper

1%

17

bioemas.com.my

Internet Source

1%

18

Mamdouh M. Ali, Fatma G. Agha. "Amelioration of streptozotocin-induced diabetes mellitus, oxidative stress and dyslipidemia in rats by tomato extract lycopene", Scandinavian Journal of Clinical and Laboratory Investigation, 2009

Publication

1%

19

Mittal, Megha, and S. J. S. Flora. "Vitamin E Supplementation Protects Oxidative Stress During Arsenic and Fluoride Antagonism in Male Mice", Drug and Chemical Toxicology, 2007.

Publication

1%

20

Submitted to Coventry University

Student Paper

1%

Exclude quotes

On

Exclude matches

< 1%

Exclude bibliography

Off