

# Turnitin Originality Report

Processed on: 27-Feb-2020 22:46 WIB  
ID: 1265287429  
Word Count: 4237  
Submitted: 1

Similarity Index	Similarity by Source
8%	Internet Sources: 4% Publications: 2% Student Papers: 7%

The Effect of Binahong Leaf (*Anredera cordifolia* [Ten] Steenis) Extract and Bay Leaf (*Eugenia polyantha* Wight) Extract Compound on Blood Glucose Level of Male Mice (*Rattus novergicus* L)  
By Diah Titik Mutiarawati

1% match (student papers from 11-May-2017)

[Submitted to Holy Name University on 2017-05-11](#)

1% match ( )

[https://pure.bond.edu.au/ws/files/25171578/Exploration\\_of\\_risk\\_factors\\_associated\\_with\\_adolescent\\_alcohol\\_consumption\\_using\\_cutting\\_edge\\_r](https://pure.bond.edu.au/ws/files/25171578/Exploration_of_risk_factors_associated_with_adolescent_alcohol_consumption_using_cutting_edge_r)

1% match (Internet from 06-Dec-2018)

<http://saspublisher.com/wp-content/uploads/2017/11/SJAMS-511D4564-4566.pdf>

1% match (student papers from 20-Sep-2013)

[Submitted to University of Reading on 2013-09-20](#)

1% match (Internet from 11-May-2018)

<http://www.mdpi.com/2072-6643/7/10/5411/htm>

1% match (Internet from 10-Nov-2019)

<https://id.scribd.com/doc/260909829/20285105-S828-Pengaruh-Campuran>

1% match (student papers from 26-Oct-2018)

[Submitted to Bowling Green State University on 2018-10-26](#)

1% match (student papers from 28-Mar-2014)

[Submitted to Curtin University of Technology on 2014-03-28](#)

1% match (publications)

[Ani Sulistyarsi, Nanda Wahyu Pribadi. "UJI AKTIVITAS ANTIBAKTERI EKSTRAK DAUN BINAHONG \(\*Anredera cordifolia\* \(ten.\) Steenis\) TERHADAP PERTUMBUHAN BAKTERI \*Staphylococcus aureus\* DAN \*Pseudomonas aeruginosa\*", \*Journal of Pharmaceutical Science and Medical Research\*, 2018](#)

1% match (publications)

[Ussama Bekkouch, Mohamed Harnafi, Ilham Touiss, Saloua Khatib, Hicham Harnafi, Chakib Alem, Souliman Amrani. "Antioxidant and Lipid-Lowering Properties of Crude Aqueous Extract and Methanolic Fraction: A Follow-Up Study", \*Evidence-Based Complementary and Alternative Medicine\*, 2019](#)

[Scholars Journal of Applied Medical Sciences \(SJAMS\) Sch. J. App. Med. Sci., 2017; 5\(11D\): 4551-4556 @Scholars Academic and Scientific Publisher \(An International Publisher for Academic and Scientific Resources\) www.saspublisher.com ISSN 2320-6691 \(Online\) ISSN 2347-954X \(Print\)](#) The [Effect of Binahong Leaf \(\*Anredera cordifolia\* \[Ten\] Steenis\)](#) Extract and Bay Leaf (*Eugenia polyantha* Wight) Extract Compound on Blood Glucose Level of Male Mice (*Rattus novergicus* L) Diah Titik Mutiarawati, Tuty Putri Sri Muljati, Indah Lestari, J. Christyaningsih Health Polytechnic Surabaya, Indonesia Abstract: The aim of this study is to explain the effect of Binahong leaf (*Anredera cordifolia* [Ten] Steenis) extract and Bay Leaf (*Eugenia polyantha* Wight) extract compound on glucose level of male mice (*Rattus novergicus* L.) blood. The mice were \*Corresponding author divided into eight groups. The normal control group was not induced by alloxan and Diah Titik Mutiarawati was not administered by Carboxy Methyl Cellulose (CMC) solution. The other groups were induced by alloxan 100 mg/kg BW in intraperitoneal manners and administered Article History by CMC (negative control group); glibenclamide (positive control group); 250 mg/kg Received: 08.11.2017 BW Binahong leaf extract (Binahong treatment group); 750 mg/kg BW bay leaf Accepted: 15.11.2017 extract (Bay Leaf treatment group); 1000 mg/ kg BW compound extract (treatment Published: 30.11.2017 extract group 1); 500 mg/kg BW (treatment extract group 2); and 250 mg/kg BW compound extract (treatment extract group 3). The administering of experimentation DOI: substances was carried out for 21 consecutive days. The result of Kruskal-Wallis and 10.21276/sjams.2017.5.11.47 single factor ANAVA ( $P < 0.05$ ) indicates that the three doses of compound significantly reduce the average glucose level on mice blood. The highest result was found on 1000 mg/kg BW dose with the normal glucose content 121.36 mg/dl on the 15th day and 85.37 mg/dl at the 22nd day. Keywords: Alloxan, *Anredera cordifolia* [Ten] Steenis, *Eugenia polyantha* Wight, Blood glucose content, *Rattus novergicus* L. INTRODUCTION Diabetes Mellitus (DM) is a disease indicated by hyperglycemia, a condition when the glucose level in the bloodstream increases exceeding the normal level. DM is the disease with second highest mortality rate on 45-54 year-old age group living in Indonesian urban areas (Depkes, 2008). Indonesia is the country with the fourth highest diabetes cases after India, China, and USA. There are some ways to reduce metabolic disorders on diabetes patients, including by organizing eating pattern and administering synthetic oral hyperglycemic agents (OHA) [1]. The excessive use of synthetic hyperglycemic medication for a long period may bring adverse effects on the patient, such as acute hypoglycemia, damages on kidneys and liver and lactic acidosis [2]. Therefore, the use of natural substances as antidiabetic agents tends to become alternative solution for the society [1]. Natural antidiabetic medications are considered more benefitted for the society due to its relatively low adverse effects compared to the synthetic medications. The administration of two or more natural medications simultaneously has proven effective because of its holistic effect in maintaining health and curing illnesses [1]. Indonesians has practiced therapy utilizing natural substances (known as polyherbal therapy) based on their empirical experiences. Tiwari and Rao [3] explain that polyherbal therapy has synergistic effect due to the substance contained within each medicinal plant. Polyherbal therapies are very beneficial because of its ability to maximize the therapeutic effects of each medicinal plant with minimum adverse effect. Several researches on the potentiality of medicinal plant extract compounds have been conducted. Agoreyo et al. [4] found that the administration of *Hibiscus sabdariffa* extract and *Zingiber officinale* extract compound potentially reduce glucose contents more optimally than the administration of single extracts. Ebong et al. [5] reported their findings indicated that the decreasing glucose contents was more optimal on the treatment

group administered by Azadirachta indica extract and Vernonia amygdalina extract compound compared to the treatment groups administered by single extracts. Among the medicinal plants known for its function as antidiabetes agents are Binahong (Madeira vine leaf) and bay leaf. These plants are potentially used as materials for polyherbal experimentation. Binahong is an ornamental plant known empirically for its medicinal use to cure kidney damage, diabetes, cardiomegaly, hematemesis, typhus, stroke, mending wounds, stomach ulcer, intestinal inflammation, dyspnea, gout, and swelling liver [6]. Bay leaf is commonly used to cure diarrhea, influenza, pneumonia, pertussis, hypertension, and diabetes [7]. The research utilizing Binahong leaf extract and bay leaf extract as antidiabetes agents has never been conducted before. The combination of these two extracts is expected to provide more optimal effect in curing diabetes. Other medicinal functions of these plants (especially Binahong, which is commonly known for its function in mending wounds) are expected to benefit diabetes patients concerning the fact that diabetes patients take much longer time to cure his/her wounds compared to non-diabetes people. The findings of preliminary research carried out by The Agency for Assessment and Application of Technology (Indonesian: Badan Pengkajian dan Penerapan Teknologi/BPPT) state that the ethanol content of Binahong leaf extract [administered at the dose of 250 mg/kg BW](#) may potentially reduce the glucose content of mice blood. Kemila [8] explains that flavonoid contents of Binahong leaf infusion may probably function as antioxidant absorbing free radicals contained within alloxan as DM inducer substance. This antioxidant activity is probably the mechanism that inhibits the function of alloxan and reduces the blood sugar contents of the alloxan-induced mice. Dewi LL et al. [10] stated that the ethanol content of bay leaf extract at the dose of 0.3-1.2 gram/kg BW might reduce the glucose level on mice blood. The dose of extract administration refers to the findings of previous research conducted by Subramanian et al. [9] and BPPT's preliminary research. The dose of compound extract administered in this study is obtained by adding the single dose of each extract, namely 250 [mg/kg BW of Binahong leaf extract](#) and 750 [mg/kg BW of bay leaf extract](#). The dose of administered compound extract is gradually decreased according to the pattern of , thus the doses varies from [1000 mg/kg BW; 500 mg/kg BW; and 250 mg/kg BW](#). Based on these backgrounds, the research question proposed in this study is how the effect of Binahong leaf extract and bay leaf extract compound administration on the glucose level of mice induced by alloxan. This study aims to identify the effect of Binahong leaf extract and bay leaf extract compound administration on [the blood glucose level of the mice](#) induced by alloxan. [The](#) writers hope that [the](#) data findings of this study can serve as preliminary information for further researches on formulation of natural extract compounds used as natural diabetes medication.

**MATERIALS AND METHODS** This experimental research applied Completely Randomized Design (CRD). The treatments were administered randomly on the 8 (eight) treatment groups with 4 (four) repetitions. The number of repetitions was determined based on Frederer formula:  $(t-1)(n-1) \geq 15$ , with t was the number of treatments and n was the number of repetitions [14]. The treatment groups of this research were: Normal Control Group (KC-1). the mice not induced by alloxan were administered by CMC 0.5% (the dose followed the body weight of each mouse) orally for 21 consecutive days. Negative Control Group (KC-2). The mice were induced by alloxan (intraperitoneal) and administered by CMC 0.5% orally for 21 consecutive days. Positive Control Group (KC-3). The mice were induced by alloxan (intraperitoneal) and administered by glibenclamide® orally (the dose 0.45 mg/kg BW orally for 21 consecutive days. Binahong Leaf Treatment Group (KPB).The mice induced by alloxan (intraperitoneal) and administered by Binahong leaf extract 250 mg/kg BW orally for 21 consecutive days Bay Leaf Treatment Group (KPS).The mice induced by alloxan (intraperitoneal) and administered by bay leaf extract 750 mg/kg BW for 21 consecutive days Extract Compound Treatment Group Dose 1 (KPD-1).The mice induced by alloxan (intraperitoneal) and administered by combined bay leaf and Binahong leaf extract compound 1000 mg/kg BW orally for 21 consecutive days. Extract Compound Treatment Group Dose 2 (KPD-2). The mice induced by alloxan (intraperitoneal) and administered by combined bay leaf and Binahong leaf extract compound 500 mg/kg BW orally for 21 consecutive days. Extract Compound Treatment Group Dose 3 (KPD-3).The mice induced by alloxan (intraperitoneal) and administered by combined bay leaf and Binahong leaf extract compound 250 mg/kg BW orally for 21 consecutive days. Diabetic Animal Model Thirty-two male mice were kept within a cage and divided into eight treatment groups (KC-1, KC-2, KC-3, KPB, KPS, KPD-1, KPD-2, and KPD-3). Each treatment group consisted of 4 (four) mice chosen randomly. Each mouse was marked using marker. The mark might be on its tails, backs, or feet [11]. The mark was used to differentiate a mouse from other mice. The mice were fed using pellets 20 gram/mice/day. Mineral water was provided as drinking water for the mice, provided in container with pipette ad libitum (unlimited). The cage was cleansed using disinfectant twice a week, rinsed, and dried up

**Extract Preparation** The extract preparation was started by washing and crushing fresh Binahong leaf and bay leaf into powder. The powdered leaves were weighed. The powdered leaves were extracted separately using ethanol solvent to produce thick extract. The thick extract was stored in refrigerator before use. **Alloxan (Diabetogen) Induction on Mice** The mice were weighed to determine the volume of alloxan administered. Alloxan was injected at the dose of 100 mg/kg BW (intraperitoneal). The volume of the injection was 0.5 ml/100 grams BW. The results of preliminary research indicated that at that dose, the injection had been enough to trigger hyperglycemic effect. Alloxan was diluted on cold aquabides at the temperature near 0 °C because alloxan was stable at that temperature. Alloxan was administered at fasting period (8-12 hours) [12]. Blood glucose level of the mice at the 0th day will be measured four days after alloxan induction. The mice were considered having diabetes if its blood glucose content exceeded 200 mg/dl. The mice with diabetes will be used as experimentation animal on this anti-diabetes examination. **CMC 0.5% Solution Preparation** 0.5 grams CMC powder was diluted on 50 ml aquadest inside 100 ml Erlenmeyer flask. The solution was homogenized on magnetic stirrer and added by aquadest until 100 ml volume. **Extract Suspension Preparation** 250 mg/kg BW Binahong leaf extract suspension was prepared by mixing 250 mg Binahong leaf with 10 ml CMC 0.5% solution in 50 ml Erlenmeyer flask. The mixture was homogenized until suspension formed. Similar procedure applied on the preparations of other extract suspension with different doses (i.e. 750 mg/kg BW bay leaf extract suspension, [1000 mg/kg BW, 500 mg/kg BW, and 250 mg/kg BW](#) compound extract suspensions). Treatment The mice were induced by alloxan (except for the normal control group) so that the mice experiencing hyperglycemia. The hyperglycemia mice were administered by experimentation plant extract for 21 consecutive days (once/day). The administration of experimentation plant extract was carried out at the same time daily (between 11 am-12 pm). The extract was administered orally using feeding tube. The normal control group (KC-1) and negative control group (KC- 2) were only administered by CMC 0.5% solution. The volume of CMC 0.5% administered depending on the body weight of each mouse. The positive control group (KC-3) was administered by glibenclamide ® solution at the dose of 0.45 mg/kg BW [8]. The treatment groups (KPB, KPS, KPD-1, KPD-2, KPD-3) were administered by extract suspensions at the determined doses [9]. The volume of administered extract suspensions was 1 ml for every 100 mg body weight, for example, the bodyweight of a mouse is 200 mg, the volume of suspension administered will be 2 ml. **Blood Sample Collection** Mice Blood samples were collected at 0th, 8th, 15th, and 22nd days of examinations. The [blood sample collection was carried out](#) after [the](#) mice were kept in fasting condition for 12 hours. The first [blood sample collection was carried out at the](#) fourth day after alloxan induction before treatment (considered as the 0th day). The second [blood sample collection was carried out at the](#) eighth day (after seven days of treatment through experimentation material administration). The third [blood sample collection was carried out at the](#) 15th day (after 14 days of treatment through experimentation material administration). The second [blood sample collection was carried out at the](#) 22nd day (after 21 days of treatment through experimentation material administration). **Blood Sample Analysis** The analysis of blood sample was carried out based on enzymatic method. **Data Processing and Analysis** Analysis on the research data was carried out based on statistical approach considering the nature of this research as experimental research. [The blood glucose level of the](#) samples taken in 0th, 8th, 15th, and 22nd days were measured using computer software. **RESULT** [Table-1: The average blood glucose levels of the](#) blood samples were taken at day 0, 7,15 and 22 of the normal control group, positive control group, negative control group, Glucose levels in Blood (mg/dL) Day 0 Day 7 Day 15 Day 22 KC-1 114.18+2.94 113.04+1.91 116.63+3.05 117.09+1.54 KC-2 233.98+21.56 245.89+16.48 226.32+24.93 276.19+2.43 KC-3 266.15+20.81 113.99+2.50 113.61+24.09 85.56+10.87 KPB 277.45+19.63 177.64+18.89 138.01+7.34 104.20+4.71 KPS 262.25+30.17 172.48+10.00 129.78+2.86 96.38+9.87 KPD-1 246.97+16.94 164.39+2.91 121.36+11.76 85.37+7.44 KPD-2 262.22+31.39 168.51+25.18 128.03+9.40 110.69+5.19 KPD-3 245.94+18.33 170.23+5.23

138.68+28.47 119.27+8.94 Note KC-1: Normal Control Group KC-2: [Negative Control Group](#) KC-3: [Positive Control Group](#) (Glibenclamide®) KPB: [Binahong Leaf Extract](#) 250 mg/kg BW KPS: Bay leaf Extract Treatment Group 750 mg/kg BW KPD-1: Compound Extract Treatment [Group Dose-1](#) (1000 [mg/kg](#) BW) KPD -2: Compound [Extract Treatment Group Dose-2](#) (500 [mg/kg](#) BW) KPD-3: Compound Extract Treatment Group Dose-3 (250 mg/kg BW) The results of Shapiro-Wilk test indicated that the glucose level of the male mice blood samples taken at the 0th day were not distributed normally with  $\alpha = 0.05$  ( $p < 0.05$ ). The results of Levene homogeneity examination indicated that the data were homogenous variance with  $\alpha = 0.05$  ( $p > 0.06$ ). The results of Kruskal-Wallis examination indicated significant differences among the treatment groups. The results of Shapiro-Wilk test indicated that the glucose level of the male mice blood samples taken at the 7th day were distributed normally with  $\alpha = 0.05$  ( $p > 0.05$ ). The results of Levene homogeneity examination indicated that the data were not homogenous variance with  $\alpha = 0.05$  ( $p < 0.06$ ). The results of Kruskal-Wallis examination indicated significant differences among the treatment groups as the effect of treatment administered. The results of Shapiro-Wilk test indicated that the glucose level of the male mice blood samples taken at the 7th day were distributed normally with  $\alpha = 0.05$  ( $p > 0.05$ ). The results of Levene homogeneity examination indicated that the data were not homogenous variance with  $\alpha = 0.05$  ( $p < 0.06$ ). The results of Kruskal-Wallis examination indicated significant differences among the treatment groups with  $\alpha = 0.05$  ( $p < 0.05$ ). The results of [Shapiro-Wilk normality test](#) indicated that [the blood glucose level of mice](#) blood taken at [the 22nd day](#) was [normally distributed](#) with  $\alpha = 0.05$  ( $p > 0.05$ ). The results of Levene homogeneity examination [indicated the data was homogeneous variance](#) with  $\alpha = 0.05$  ( $p < 0.05$ ). the results of single factor ANOVA examination indicated that the treatment administered had affected the blood glucose level at the 22nd day as indicated by significant differences among the treatment group  $\alpha = 0.05$  ( $p > 0.05$ ). There are significant differences among the treatment groups: KC-1, KC-3, KPB, KPS, KPD-1, KPD-2, and KPD-3 compared to KC-2, KPD-3 compared to KC-1, KPD-3 compared to KPS, KPD-2 and KPD-3 compared to KPD-1; and KPD-2 and KPD-3 compared to KPD-1. DISCUSSION Data on blood glucose level at the 0th day of this research was needed to determine the uniformity [of blood glucose level of the](#) experimentation animals used in this research. Hyperglycemia condition found on the experimentation animal is the result of alloxan induction at the dose of 100 mg/kg BW. The determination of the alloxan dose used in this research was based on initial orientation. Alloxan is one of diabetogen (substance that causes diabetes) that is selective in causing damages on Langerhans  $\beta$ -cells of pancreas. Szkudelski [15] state that alloxan induction may trigger permanent hyperglycemia condition in quick time (for about 2-3 days). The examination of [the blood glucose](#) contents [of the](#) male [mice](#) blood samples confirmed this statement. The blood samples of the 32 male mice indicated that the mice were at hyperglycemic condition 4 days after alloxan induction. The normal control group (KC-1) was not administered by experimentation extract. However, the group was administered by Carboxy Methyl Cellulose (CMC) 0.5% during the treatment period, similar to the other treatment groups in order to prevent research bias. The administration of CMC 0.5% on KC-1 also aims to prove that CMC 0.5% is not the factor that affects blood glucose level of *Rattus novergicus* L. Murray et al. [2] stated that CMC could not be digested by mammals because mammals did not have certain enzyme to carry out hydrolysis on the cellulose. Therefore, the blood glucose content is not affected by CMC. The average blood glucose contents of *Rattus novergicus* L. induced by alloxan at the day 0 was ranged from 233.98  $\pm$  21.56 mg/dl to 277.65  $\pm$  19.63 mg/dl while the average blood level content of the normal control group was 114.18  $\pm$  2.94 mg/dl. This condition confirms the theory stating that the normal glucose blood level of *Rattus novergicus* L. was 50-135 mg/100 ml [13]. Based on the blood glucose levels of the samples collected at day 0, it can be concluded that the blood glucose level of *Rattus novergicus* L. after alloxan induction tends to be uniform. This finding is supported by the results of Kruskal-Wallis non- parametric test with  $\alpha = 0.05$  ( $p > 0.05$ ) indicating there is no significant difference among the treatment group. The data on *Rattus novergicus* L. blood glucose content of the normal control group (KC-1) indicates that there is no blood glucose level exceeds 200 mg/dl. The administration of compound extract for 14 consecutive days shows significant effect on the decreasing of blood glucose content of *Rattus novergicus*. This condition can be found on Table 3 and the results of Kruskal Wallis test ( $p < 0.05$ ,  $\alpha = 0.05$ ). the administration of compound extract in various doses (1000, 500, and 250 mg/dl) for 14 consecutive days has proven decreasing the blood glucose level of *Rattus novergicus* L. induced by alloxan. Data on blood glucose content of the samples taken at 15th day indicated that average levels of blood glucose on the KPD-1 and KPD-2 was lower than KPB and KPS. This finding proves that the combined (compound) extract seems to be more effective in reducing blood glucose level compared to the single extracts. The decreasing dose of compound extract (as seen on KPD-2) turns out to provide greater results compared to KPB and KPS. Mutual interactions between bioactive substances contained within each medicinal plant might be the factor explaining why the compound extract is more effective in reducing [blood glucose level](#). Based on [the data of blood glucose](#) level of the experimentation animals taken at the 22nd day of examination (as presented on Table 3.4 above) and the results of ANOVA test with  $\alpha = 0.05$  ( $p < 0.05$ ) (see Appendix), it is found that there was significant differences of the average blood glucose content on KPD-3 compared to KC-1; KPD-3 compared to KPS; KPD-2 and KPD-3 compared to KC-3; KPD-2 and KPD-3 to KC-1; KPD-1 compared to KPD-2 and KC to KPD-3. The decreasing of compound extract dose still potentially reduced the blood glucose content although the average blood glucose level of KPD-2 and KPD-3 at the 22nd day seemed to be higher than single extracts (KPB and KPS). The dose of each extract combined as KPD-2 and KPD-3 may contain a number of active compounds that reduce blood glucose content lower than KPD-3. Although the average blood glucose levels in KPD-3 = 119.27  $\pm$  8.94 mg / dl is still higher than KC3 = 85.56  $\pm$  10,87mg / dl and KC1 = 117.09  $\pm$  1.54 mg / dl, KPD- 3 is capable of approaching the average value of blood glucose levels in a single extract group (i.e. the single extract KPB = 104.20  $\pm$  4,71mg / dl and KPS = 96.38  $\pm$  9,87 mg / dl). It shows that the smallest dose on the compound extract (KPD-3) is effectively able to lower blood glucose level better. This is consistent with the theory that states the benefits of polyherbal therapy because it can increase the ability of therapeutic doses and side effects as small as possible [3]. CONCLUSION AND RECOMMENDATION Based on the findings of this study examining the effect of combined (compound of) Binahong leaf and bay leaf extract in reducing blood glucose level of male mice (*Rattus novergicus* L.) induced by alloxan, the conclusion of this study are: The administration of Binahong (*Anredera cordifolia* [Ten] Steenis) leaf extract and bay leaf (*Eugenia polyantha* Wight) compound extract at the doses of 1000 mg/kg BW, 500 mg/kg BW, and 250 mg/kg BW is able to lower the blood glucose levels of male mice (*Rattus novergicus* L.) induced by alloxan. Further researchers should focus their studies on toxicity effect of Binahong (*Anredera cordifolia* [Ten] Steenis) and bay leaf (*Eugenia polyantha* Wight) of the mice with diabetes and the impacts of the extract administration on internal organs such as kidneys, liver, and pancreas. These studies need to be conducted in order to gain more comprehensive information on the benefits and safety of Binahong leaf extract and bay leaf extract as natural treatment for diabetes mellitus. REFERENCES 1. Tjay TH, Rahardja K. Obat-obat penta: khasiat, penggunaan dan efek-efek sampingnya. Elex Media Komputindo; 2007. 2. Murray RK, Granner DK, Mayes PA, Rowell VW. Biokimia harper. Ed.25, Terj. dari Harper's Biochemistry, oleh Andry H. Penerbit Buku Kedokteran EGC, Jakarta: ix + 883 hlm. 2003. 3. Tiwari AK, Rao JM. Diabetes mellitus and multiple therapeutic approaches of phytochemicals: Present status and future prospects. Current science. 2002 Jul 10;30-8. 4. Agoreyo FO, Agoreyo BO, Onuorah MN. Effect of aqueous extracts of *Hibiscus sabdariffa* and *Zingiber officinale* on blood cholesterol and glucose levels of rats. African Journal of Biotechnology. 2008;7(21). 5. Ebong PE, Atangwho IJ, Eyong EU, Egbung GE. The antidiabetic efficacy of combined extracts from two continental plants: *Azadirachta indica* (A. Juss)(Neem) and *Vernonia amygdalina* (Del.)(African bitter leaf). American Journal of Biochemistry and Biotechnology. 2008;4(3):239- 44. 6. Manoi F. Binahong (*Anredera cordifolia*) Sebagai Obat. Warta Penelitian dan Pengembangan Tanaman Industri. 2009 Apr;15(1):3-5. 7. Hariana A, Obat T. Khasiatnya, Seri 3. Jakarta: Penebar Swadaya. 2006:160-1. 8. Kemila M. Uji Aktivitas Antidiabetes Mellitus Ingus Daun Binahong Pada Tikus Jantan. Skripsi jurusan Farmasi FMIPA-UII, Yogyakarta:iv +55 hlm. 2010. 9. Subramanian R, Asmawi MZ, Sadikun A. Effect of andrographolide and ethanol extract of *Andrographis paniculata* on liver glycolytic, gluconeogenic, and lipogenic enzymes in a type 2 diabetic rat model. Pharmaceutical biology. 2008 Jan 1;46(10-11):772-80. 10. Dewi IL. Uji Aktivitas Antidiabetes Ekstrak Etanol Daun Salam (*Eugenia polyantha*) terhadap tikus galur wistar yang diinduksi aloksan (Doctoral dissertation, Universitas Muhammadiyah Surakarta).2013. 11. Radji M. Harmita. 2004. Buku Ajar Analisis Hayati.:47-55. 12. Fröde TS, Medeiros YS. Animal models to

test drugs with potential antidiabetic activity. *Journal of Ethnopharmacology*. 2008 Jan 17;115(2):173-83. 13. Carvalho EN, Carvalho NA, Ferreira LM. Experimental model of induction of diabetes mellitus in rats. *Acta Cirurgica Brasileira*. 2003;18(SPE):60-4. 14. Hanafiah AK. Rancangan percobaan: Teori dan Aplikasi. Ed.ke2. Penerbit PT. Raja Grafindo Persada, Jakarta :xii +238 hlm. 1997. 15. Johnson C, Ahsan N, Gonwa T, Halloran P, Stegall M, Hardy M, Metzger R, Shield III C, Rocher L, Scandling J, Sorensen J. Randomized trial of tacrolimus (prograf) in combination with azathioprine or mycophenolate mofetil versus cyclosporine (neoral) with mycophenolate mofetil after cadaveric kidney transplantation1, 2. *Transplantation*. 2000 Mar 15;69(5):834-41. Diah Titik Mutiarawati [et al., Sch. J. App. Med. Sci., Nov 2017; 5\( 11D\):4551-4556](#) Diah Titik Mutiarawati [et al., Sch. J. App. Med. Sci., Nov 2017; 5\( 11D\):4551-4556](#) Diah Titik Mutiarawati [et al., Sch. J. App. Med. Sci., Nov 2017; 5\( 11D\):4551-4556](#) Diah Titik Mutiarawati [et al., Sch. J. App. Med. Sci., Nov 2017; 5\(11D\):4551-4556](#) [Available online at http://saspublisher.com/sjams/](http://saspublisher.com/sjams/) 4551 [Available online at http://saspublisher.com/sjams/](http://saspublisher.com/sjams/) 4552 [Available online at http://saspublisher.com/sjams/](http://saspublisher.com/sjams/) 4553 [Available online at http://saspublisher.com/sjams/](http://saspublisher.com/sjams/) 4554 Available online at <http://saspublisher.com/sjams/> 4555 Available online at <http://saspublisher.com/sjams/> 4556