

Turnitin Originality Report

Processed on: 05-Aug-2020 20:25 WIB
 ID: 1366198464
 Word Count: 3062
 Submitted: 1

Similarity Index

12%

Similarity by Source

Internet Sources:	8%
Publications:	1%
Student Papers:	5%

Factor Analysis on Measles
 after The 2017 Measles
 Rubella Vaccine Campaign in
 Magetan District By Tutiek
 Herlina

4% match (Internet from 18-May-2019)

http://ijshr.com/IJSHR_Vol.4_Issue.1_Jan2019/IJSHR0032.pdf

2% match (student papers from 01-Apr-2020)

[Submitted to Aspen University on 2020-04-01](#)

2% match (Internet from 29-May-2019)

<http://journal.aloha.academy/index.php/ajjha/article/download/54/pdf>

2% match (Internet from 04-Feb-2020)

<http://heanoti.com/index.php/hn/article/download/hn20110/87>

1% match (student papers from 04-Jan-2019)

[Submitted to iGroup on 2019-01-04](#)

1% match (student papers from 19-Mar-2012)

[Submitted to University of Leeds on 2012-03-19](#)

Factor Analysis on Measles after The 2017 Measles Rubella Vaccine Campaign in Magetan District Tutiek Herlina¹ (corresponding author), Triana Septianti Purwanto² ¹Health Polytechnic of Surabaya, Indonesia (herlinatutiek97@gmail.com) ²Health Polytechnic of Surabaya, Indonesia Submitted: January 22, 2019 -Revised: April 1, 2019 -Accepted: April 28, 2019 -Published: April 30, 2019 ABSTRACT Measles is an infectious disease, infecting many children and becomes a major cause of child mortality in Indonesia. This disease is highly contagious, transmitted by viruses through droplets. Undeveloped immunity and low immunization coverage make it potentially epidemic. According to the 2016 Health Profile of Magetan Regency, measles cases were ranked first. Indonesia is committed to achieve the elimination of measles in 2020, with strategies such as the Measles Rubella vaccine campaign for children between 9 months-15 years. The research purpose was to analyze the factor of measles after the 2017 measles rubella vaccine campaign in Magetan Regency. This research used retrospective case control research design, the case sample was all the measles patients after the MR vaccine campaign until July 2018, as many as 16. The control sample was a part of the Magetan Regency population who did not suffer measles, as many as 64, determined with a ratio of 1:4. The research variables were immunization status, gender, exclusive breastfeeding, patient contact history, and knowledge of measles. The results [showed that there is an effect of immunization status on the incidence of measles. The probability of someone who was not immunized is 81% for measles infection compared to those who are immunized. There is no influence of gender,](#)

exclusive breastfeeding, patient contact history, and knowledge of measles on the incidence of measles. Conclusion: immunization factor is the dominant factor that influence the incidence of measles. Suggestion: to reduce morbidity and mortality rates due to measles, Health Center (Puskesmas) is expected to increase the coverage of measles immunization, supported by highest quality of vaccine and techniques.

Keywords: measles, campaign, rubella measles vaccine

INTRODUCTION

Background Measles is an infectious disease which infects many children and is a major cause of child mortality in Indonesia. This disease is highly contagious, transmitted by viruses through droplets when coughing or sneezing. Measles is a potential epidemic if the immunity is not developed and the immunization coverage is low(1). Measles cases in Indonesia increase from 2013 to 2015. Surveillance activities reported that more than 11.000 cases of suspected measles with laboratory confirmation results of 12-39% are definite measles. The number of cases from 2010 to 2015 is estimated to have 23.164 cases. This number is estimated to be lower than the actual number in the field, considering there are still many unreported cases, especially from private services and the completeness of the surveillance reports that are still low(2). According to the 2016 Health Profile of Magetan Regency, measles cases were ranked first(3). The measles immunization status affects the protection from measles. Population density is a factor which influences the spread of the disease, because dense settlements are fertile nurseries for viruses and facilitate transmission. Decreased vitamin A levels lead to increased child mortality(4). Measles can cause complications including Bronchopneumonia, Acute Otitis Media, Encephalitis and some children can suffer Enteritis(5). The Global Vaccine Action Plan (GVAP) of measles is targeted to be eliminated in 5 WHO regions by 2020. The 2012-2020 Global Measles & Rubella Strategic Plan maps out the strategies needed to reach the target of world without measles, rubella or CRS. One of the five strategies is achieving and maintaining a high level of immunity by giving two doses of vaccine containing measles through routine and additional immunization with high coverage (>95%) and evenly distributed. Indonesia has committed to achieve an elimination of measles and CRS control by 2020, with strategies such as routine strengthening immunization to achieve measles coverage of $\geq 95\%$ evenly at all levels; implementing measles crash program in children aged 9- 59 months in 185 districts/cities in August-September 2016; making MR (Measles Rubella) vaccine campaign for children aged 9 months-15 years in stages in 2 phases of August-September 2017 and 2018; introduction of MR vaccines to routine immunization programs in October 2017 and 2018; conducting case-based rubella measles and CRS sentinel surveillance in 13 hospitals; and fully investigating the Extraordinary Phenomenon of measles outbreak(2). Goal This research to analyzed the influence of immunization, gender, exclusive breastfeeding, contact history with patients, and knowledge of measles toward the incidence of measles after the 2017 measles rubella vaccine campaign in Magetan Regency.

Hypothesis Immunization status influenced to incidence of measles; gender influenced to incidence of measles; exclusive breastfeeding influenced to incidence measles; contact with measles sufferers influenced to incidence of measles; knowledge about measles influenced to incidence measles.

METHODS

This research used retrospective case control design. The sample cases were all measles patients after the 2017 MR vaccine campaign, which are as many as 16 people. The control group sample was a part of the Magetan Regency population who did not suffer measles, as many as 64 people. It was taken with consecutive techniques with a ratio of cases and controls 1:4 because the prevalence of measles cases was unknown(6). The research variables were immunization, gender, exclusive breastfeeding, contact history with patients, and knowledge of measles. The instrument of data collection was a questionnaire. The data analysis used Chi Square statistical test and logistic regression. This research had

received ethical approval. RESULTS Relation between Immunization Status, Gender, Exclusive Breastfeeding, Patient Contact History, and Knowledge of Measles with The Incidence of Measles The relation between immunization status, gender, exclusive breastfeeding, contact history and knowledge of measles with the incidence of measles [can be seen in the following table. Table 1](#). The relation between immunization status, gender, exclusive breastfeeding, patient contact history, knowledge [of measles](#) and [the incidence of measles](#) Variables Infected f % [Measles Not](#) infected f % [Total](#) f % [p-value](#) Immunization status 0.000 Not immunized 9 7 16 81.8 10.1 20.0 2 62 64 18.2 89.9 80.0 11 69 80 100 100 100 Gender 0.014 Men women total 13 3 16 30.2 8.1 20.0 30 34 64 69.8 91.9 80.0 43 37 80 100 100 100 Exclusive breastfeeding history 0.214 Not exclusive Exclusive Total 7 9 16 15.2 26.5 20.0 39 25 64 84.8 73.5 80.0 46 34 16 100 100 100 Contact history with patient 0.029 Presence absence total 10 6 16 32.3 12.2 20.0 21 43 64 67.7 87.3 80.0 31 49 80 100 100 100 Knowledge of measles 0.138 Less well Total 9 7 16 28.1 14.6 20.0 23 41 64 71.9 85.4 80.0 32 48 80 100 100 100 Most of measles the patients who do not get immunizations are male, get breastfed exclusively, have a history of contact with measles patients, but have good knowledge about measles. Table 2. The effect of immunization status, gender, exclusive breastfeeding, patient contact history, knowledge of measles on the incidence of measles Variables B Significance Exp (B) Under KI 95% Above sex(1) 1.312 0.106 3.713 0.758 18.186 immunization(1) 3.299 0.001 27.077 4.105 178.600 Stage 1a contact(1) 0.722 0.333 2.058 0.477 8.873 knowledge(1) 0.298 0.686 1.348 0.317 5.730 exclusive breastfeeding(1) 0.197 0.800 1.217 0.267 5.560 Constant -3.504 0.001 0.030 sex(1) 1.304 0.107 3.684 0.754 17.996 immunization(1) 3.240 0.000 25.535 4.120 158.262 Stage 2a contact(1) 0.719 0.335 2.051 0.476 8.841 knowledge(1) 0.302 0.682 1.352 0.319 5.735 Constant -3.370 0.000 0.034 sex(1) 1.316 0.104 3.728 0.763 18.204 Stage 3a immunization(1) 3.293 0.000 26.917 4.378 165.500 contact(1) 0.731 0.326 2.078 0.483 8.934 Constant -3.260 0.000 0.038 Stage 4a sex(1) 1.259 0.120 3.521 0.721 17.198 immunization(1) 3.504 0.000 33.262 5.711 193.739 Constant -2.955 0.000 0.052 Stage 5a immunization(1) 3.685 0.000 39.857 7.137 222.577 Constant -2.181 0.000 0.113 The statistical results through five stages showed that immunization status variable has a dominant influence on the incidence of measles, with a value of $p=0.000$ ($p<0.05$). The regression equation model obtained: $Y = -2,181 + (3,685 \times 1) = 1,504$ $P = 1 / (1+0,222) = 0,81$ or 81%. The incidence of measles probability in people who are not immunized is 81%. DISCUSSION Age The ages of measles patients after the 2017 MR vaccine campaign in Magetan Regency are as follows. The youngest was 3 years old, the oldest was 16 years old and more in primary school age, 6-12 years old. All ages had the same susceptibility to infection(1). The results showed that the most cases of measles were in 30 months of age(7), then adults, followed by children aged 8 months-5 years, and the least is under 8 months of age(8). The highest incidence of measles is in the age group <5 years(9). The three results of the research showed distribution in different age groups. The spread of age distribution on measles patients is the basis of MR vaccine campaign given to children aged [9 months to 15 years, with a high coverage \(minimum 95%\) and evenly](#) distributed, thus is [expected to form immunity so as to reduce the virus's transmission to a more mature age and protect the group when entering the reproductive age](#). Residence Measles patients after the 2017 MR vaccine campaign in Magetan District reside in Karas, Magetan, Parang, Ngariboyo, Kartoharjo, Bendo, Plaosan and Kawedanan Sub-districts. Measles is related to population density and immunization coverage. The incidence of measles tendency is lower in an isolated population and with a small population of <400.000 people, as well as regions with high immunization coverage(1). The results showed that population density is a risk factor of measles(10),(11). According to the

2017 Health Office Profile of Magetan Regency, Karas, Bendo and Parang Sub-districts are not included in the criteria for high population density, but in terms of immunization coverage they are still below the target. Even though Magetan, Kawedanan and Kartoharjo Sub-districts are included in high population densities, but the immunization coverage is high. Whereas the case in Ngariboyo Sub-district, besides the population density is high, the immunization coverage is also low. Immunization Status There is an influence of immunization on the incidence of measles. The tendency of individuals who did not get immunization to be infected by measles is 81% compared to individuals who got immunization. Measles patients after the 2017 MR vaccine campaign in Magetan Regency, most of them did not get immunization. The presence of immunization is expected to protect people from infection and to have important effects in the disease epidemiology by changing the age distribution. Giving immunization to babies will reduce the infection agent's transmission and reduce the chance of someone who is vulnerable to exposure. Giving immunization to adolescence will prevent the incidence of measles in the reproductive age(2). Supporting research results including Yu Wen Zhou's, Shiu Tie Jun's research on the analysis of epidemiological characters in China from 2004 to 2006 concluded that there was a decrease in cases after measles immunization campaigns in Xinjian and Guizhov(12). Research by Mujiati, Muntahar, Rakmiwati on measles risk factors concluded that immunization status was the cause of measles incidence in children aged 1-14 years in Metro Pusat Sub-district, Lampung Province(13). Gender There is no influence of gender on the incidence of measles. Measles patients after the 2017 MR vaccine campaign in Magetan Regency were mostly men. Women had higher antibody titers than men, this condition will affect immunity so infection is higher in men, as well as the risk of measles infection(1). The results of research by Yahya Mohammed, that the ratio of men and women to measles infection is 2:1. The absence of gender influence on the incidence of measles, this is likely because there are more dominant factors(7). Because male is one of the factors that contributes to measles, parents should try to increase the immunity of their male babies, children, and adolescents. Exclusive breastfeeding There is no influence of exclusive breastfeeding history on the incidence of measles. Measles patients after the 2017 MR vaccine campaign in Magetan District are more with a history of obtaining exclusive breastfeeding. The results of this research do not support the results of Ardiyanto, Kirwono, Kusumawati's research which shows that there is a relation between exclusive breastfeeding and measles. Exclusive breastfeeding is breast milk given to babies up to the age of 6 months without other foods. There are more than 30 types of immunoglobulin contained in breast milk, especially Ig A and Ig G, which protect the baby's body against infection(1). Even having breastfed exclusively, but more cases of measles possibly due to reduced immunity because the patient is 3 years old or older. Contact Patient History There is no influence of contact history with patient on the incidence of measles. Measles patients after the 2017 MR vaccine campaign in Magetan Regency, most of them have a history of having contact with measles patients before. When someone is exposed to measles, 90% of people who interact closely with patients can get infected if they are not immune to measles(2). Even though there is contact with the patient, as long as the body's immunity is good, they can avoid the possibility of contracting measles. Knowledge of Measles There is no influence of knowledge about measles on the incidence of measles. Measles patients after the 2017 MR vaccine campaign in Magetan Regency mostly have lack of knowledge about measles. Knowledge or cognitive is very important domain in shaping a person's actions (overt behavior). Based on the experience and the study, behaviors that are by knowing the nature of infectious diseases, modes of transmission, methods of prevention, handling methods, and factors that support measles to easily infect, it will support someone to act

or do things that can prevent transmission and contract disease.

CONCLUSION Measles immunization affects a person's susceptibility to measles. The probability of measles incidence in individuals who are not immunized is 81% compared to those who get immunization. Gender, exclusive breastfeeding, contact history with patient and knowledge of measles do not affect a person to get measles infection. To reduce morbidity and mortality due to measles, the health center (puskesmas) is expected to increase the coverage of measles immunization with the support of highest quality of vaccine and techniques. People who are the target of the measles immunization program are expected to participate in the program. . REFERENCES 1. Setiawan IM. Measles (Penyakit Campak). Jakarta: PT Sagung Seto; 2008. 2. Kemenkes RI. Guideline of Technical Campaign and Introduction Measles Rubella (MR) Immunization (Petunjuk teknis Kampanye dan Introduksi Imunisasi Measles Rubella (MR)). Jakarta: Kemenkes RI; 2017. 3. Dinkes Kab. Magetan. Health Profile of Magetan District for 2016 (Profil Kesehatan Kabupaten Magetan 2016). Magetan: Dinkes Kab. Magetan; 2017. 4. Sudfeld CR; Navar AM; Halsey NA. Effectiveness of Measles Vaccination and Vitamin A treatment. Int J Epidemiol. 2010;39(Suppl_1):48-55. 5. Nelson. Children Health Science (Ilmu kesehatan Anak). 15th edition. Jakarta: EGC; 2000. 6. Murti B. Principle and Methods of Epidemiology Research (Prinsip dan Metode Riset Epidemiologi). Yogyakarta: Gadjah Mada University Press; 1995. 7. Yahya M. Descriptive Analysis of Measles Cases Seen in A tertiary Health Facility Sokoto, North-Western Nigeria-Implication of Disease Eradication. Glob J Med Public Heal. 2017;6(2). 8. Ying, NIU; Chun-mei, CHAO; Lan Y. Epidemiology and Clinical Analysis on 95 Cases of Measles. Chinese J Infect Control. 2007;2. 9. Nurani, DS; Ginanjar, P; Dian S. Descriptive Epidemiological Measles at Cirebon City for 2004-2011 (Gambaran Epidemiologi Kasus Campak Di Kota Cirebon tahun 2004-2011). J Kesehat Masy FKM UNDIP. 2012;1(2):293-304. 10. Giarsawan, N; Asmara, Suarta; Yulianti A. Influencing Factors for Measles at Tejakula Buleleng District's Public Health Center for 2012 (Faktor-Faktor Yang Mempengaruhi Kejadian Campak di Wilayah Puskesmas Tejakula I Kabupaten Buleleng tahun 2012). J Kesehat Lingkung. 2014;4(2):140-5. 11. Isu, AL; Weraman PP. Spatial Analyzis of Measles Outbreak by Geographical Information System (Kajian Spasial Faktor Risiko Terjadinya kejadian Luar Biasa Campak Dengan Geographical Information System). J media Kesehat Masy Indones. 2015;12(4). 12. YU, Wen-zhou; SHIU T. Analysis on Epidemilological Characteristics And Control Measures of Measles in China During 2004-2006. Chinese J Vaccine Immunization. 2007;2. 13. Mujiati, E; Mutahar, R; Rahmawati A. Risk Factors of Measles Cases on Children Aged 1-14 year at MetroPusat Sub district, Lampung Province for 2103-2014 (Faktor Risiko Kejadian Campak Pada Anak Usia 1-14 tahun di Kecamatan Metro Pusat Propinsi Lampung tahun 2013-2014). J Ilmu Kesehat Masy. 2015;6(2):100-12. [Aloha International Journal of Health Advancement](http://journal.aloha.academy/index.php/aijha) (AIJHA) Volume 2 Number 4, April 2019 <http://journal.aloha.academy/index.php/aijha> ISSN 2621-8224 RESEARCH DOI: <http://dx.doi.org/10.33846/aijha20404> [Aloha International Journal of Health Advancement](http://journal.aloha.academy/index.php/aijha) (AIJHA) Volume 2 Number 4, April 2019 <http://journal.aloha.academy/index.php/aijha> ISSN 2621-8224 RESEARCH DOI: <http://dx.doi.org/10.33846/aijha20404> [Aloha International Journal of Health Advancement](http://journal.aloha.academy/index.php/aijha) (AIJHA) Volume 2 Number 4, April 2019 <http://journal.aloha.academy/index.php/aijha> ISSN 2621-8224 RESEARCH DOI: <http://dx.doi.org/10.33846/aijha20404> [Aloha International Journal of Health Advancement](http://journal.aloha.academy/index.php/aijha) (AIJHA) Volume 2 Number 4, April 2019 <http://journal.aloha.academy/index.php/aijha> ISSN 2621-8224 RESEARCH DOI: <http://dx.doi.org/10.33846/aijha20404> 92 | Publisher: Alliance of Health Activists (AloHA) 93 | Publisher: Alliance of Health Activists (AloHA)

94 | Publisher: Alliance of Health Activists (AloHA) 95 | Publisher: Alliance of Health Activists (AloHA) 96 | Publisher: Alliance of Health Activists (AloHA)