

# Turnitin Originality Report

Processed on: 01-Oct-2020 9:01 PM CDT  
 ID: 1393843933  
 Word Count: 3484  
 Submitted: 2

Similarity Index

21%

## Similarity by Source

Internet Sources: 21%  
 Publications: 3%  
 Student Papers: 4%

Potency Of Ginger Extract  
 (Zingerber Officinale Roscoe)

As Antioxidant And

Hepatoprotector For Liver And

Pulmonary Tissue In Exposed  
 Mice (Musculus), 1. Umi

Rahayu, 2. Imam Thohari 3.

Ngadino By Heruswn Forikes

17% match (Internet from 16-Aug-2020)

<https://journalcra.com/article/potency-ginger-extract-zingerber-officinale-roscoe-antioxidant-and-hepatoprotector-liver-and>

2% match (student papers from 24-Jun-2020)

[Submitted to Universitas Nahdlatul Ulama Surabaya on 2020-06-24](#)

1% match (Internet from 27-Jul-2020)

<https://cybermedlife.eu/index.php/scientific-studies-biomedical-literature/pharmacological-actions/giulio-tarro>

1% match (student papers from 24-Jun-2020)

[Submitted to Universitas Nahdlatul Ulama Surabaya on 2020-06-24](#)

1% match (Internet from 17-Apr-2020)

<https://es.scribd.com/document/318854508/3>

1% match (student papers from 11-Aug-2020)

[Submitted to auf on 2020-08-11](#)

< 1% match (student papers from 20-Jan-2014)

[Submitted to Universitas Muhammadiyah Surakarta on 2014-01-20](#)

< 1% match (publications)

[Hana Nika Rustia, Bambang Wispriyono, Dewi Susanna, Fitra N. Luthfiah. "Organophosphate Pesticide Exposure Effects toward Inhibition of Blood Cholinesterase Activity among Vegetable Farmers. Organophosphate pesticides can inhibit blood cholinesterase in human body", Makara Journal of Health Research, 2011](#)

[Available online at http://www.journalcra.com](http://www.journalcra.com) INTERNATIONAL JOURNAL OF CURRENT RESEARCH International Journal of Current Research Vol. 12, Issue, 01, pp. 9281-9284, January, 2020 DOI:

<https://doi.org/10.24941/ijcr.37708.01.2020> ISSN: 0975-833X

RESEARCH ARTICLE [POTENCY OF GINGER EXTRACT \(ZINGERBER OFFICINALE ROSCOE\) AS ANTIOXIDANT AND HEPATOPROTECTOR FOR LIVER AND PULMONARY TISSUE IN EXPOSED MICE \(MUS MUSCULUS\) BY ORGANOPHOSPHATE PESTICIDES \\*Umi Rahayu, SKM. M., Kes, Imam Thohari ST, MMKes, Ngadino, S. Si., M.Psi](#) Health Polytechnic of Ministry of Health in Surabaya [ARTICLE INFO Article History: Received 14th October, 2019 Received in revised form 28th November, 2019 Accepted 29th December, 2019 Published online 30th January, 2020](#) Key Words: Pesticides, Ginger, Liver and Pulmonary. [ABSTRACT Pesticides as bioactive](#)

substances contain poison for killing plant-disturbing organism. Pesticides exposure can cause damage for liver and pulmonary tissue. Each active substance that is contained in pesticides causes different symptoms of poisoning. The symptoms of poisoning (subjective complaints) which are caused as the impact of pesticides will be improved by us through providing ginger extract. Moreover, this research will be applied to mice (Mus musculus). This research aimed at analyzing the influence of providing ginger extract (Zingiber officinale Roscoe) as antioxidant against the protection of liver and pulmonary damage due to pesticides exposure in mice. Design of this research was Quasi Experimental research by using simple experimental design (Post Test Only Control Group Design). Sample in this research was healthy male Swiss Webster white mice. An examination was conducted in order to investigate tissue damage on experimental animals' liver and pulmonary. The research result showed that there was liver cell damage in K0 group (without any treatment). Besides, there were normal cell and the lightest damage (less than 25%). In K1 group, it was occurred cell degeneration and hepatocytic damage in mild, moderate, and chronic hepatocyte damage. In K3 group, it was occurred cell degeneration and hepatocyte damage in mild, moderate, and chronic level. However, improving liver cells was through providing ginger extract. Through adding ginger extract, improving damage on pulmonary organ cells was also much better. In conclusion, by providing ginger extract in mice which were exposed by pesticides had no role in repairing liver cell damage. In contrary, in pulmonary organ, it was occurred an improvement of mice's pulmonary organ cell. Nevertheless, for further research, it needs to be conducted a research in smaller dose of ginger extract. Copyright © 2020, Umi Rahayu [et al.](#) [This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.](#) Citation: Umi Rahayu, SKM. M. Kes, Imam Thohari ST, MMKes, Ngadino, S.Si., M.Psi, 2020. "Potency of ginger extract (zingerber officinale roscoe) as antioxidant and hepatoprotector for liver and pulmonary tissue in exposed mice (mus musculus) by organophosphate pesticides", *International Journal of Current Research*, 12, (01), 9281-9284.

INTRODUCTION The use of organophosphate pesticides in Indonesia is conducted by spraying after the attack of pests and this is worsened by unconcern from the farmers regarding the use of organophosphates which is dangerous for bodily health. Farmers in Indonesia need pesticides in their life, but they are less care regarding the impact of pesticides. The symptoms of being poisoned (subjective complaint) from organophosphate and carbamate groups are such as arising certain muscle movements, blurred vision, watery eyes, foaming mouth, sweating much, salivating much, dizziness, nausea, convulsions, vomiting, fast heart, shortness of breath, and rapid heartbeat. \*Corresponding author: Umi Rahayu, SKM. M..Kes, Health Polytechnic of Ministry of Health in Surabaya. Several substances which are contained in pesticides (such as organophosphate and carbamate groups) can reduce the ability of cholinesterase enzyme to hydrolyse acetylcholine. Thus, rate of stimulation delivering in nerve impulses is hindered and finally, it will cause abnormality of nervous system (Rasyid, 1995). Furthermore, main function of liver is detoxification of toxic substances in the body. Liver disfunction will impact on the hindered formation of red blood cells and the disruption of food substances metabolism. Liver is an organ that can metabolize and excrete several chemicals (Mansur, 2008). In order to increase the improvement of liver function, particularly for farmer who is exposed by pesticide poison, it needs to be looked for natural substances which are easy to be obtained, cheap, easy to be processed, and non-toxic. Ginger is one of the most used spices in society. It contains antioxidant- gingerol as food ingredient and Indonesian traditional beverage / medicine. Umi Rahayu et al. Potency of ginger extract (zingerber officinale roscoe) as antioxidant and

[hepatoprotector for liver and pulmonary tissue in exposed mice \(mus musculus\) by organophosphate pesticides](#) Moreover, it has been conducted a research against SGPT (Serum Glutamic Pyruvate Transaminase) level and description of hepatic histopathology due to exposure of Allaetrin by Achmad Fathir in 2010. Rachmaniyah et.al, in 2018 with the title "Pengaruh rimpang Jahe (Zingerber officinale Roscoe) terhadap aktivitas enzim Kolinestarase akibat paparan pestisida pada mencit terpapar pestisida Organofosfat" (Effect of Ginger rhizome (Zingerber officinale Roscoe) on cholinestarase enzyme activity due to pesticide exposure in exposed mice by Organophosphate pesticides) has proved that ginger has very strong antioxidant activity and can improve cholinesterase enzyme activity in mice which are exposed by pesticides. In conducted research by Fugio et al regarding the characteristic of antioxidant in chemical component of ginger was found the component of several compounds which had important role in antioxidant activity of ginger: [gingerdiol, gingerol, caffeic acid, camphene, capsaicin, chlorogenic acid, curcumin, delphinidin, eugenol, ferulic acid, gamma](#) terpinen, [isoeugenol, melatonin, myrcene](#), sam vanilat, [vanillin](#), and [zingerone](#). (Utami <sup>Δ</sup> Meryalita Prihatin et al, 2012). Conducted research by Yuli Haryani, 2016, also proved that ginger rhizome water extract (Zingiber off Cinale ) could reduce AgNO<sub>3</sub> in simple biosynthesis of silver nanoparticles at room temperature simply. Result of previous research (Imam Thohari et al, 2017) explained the function of red guava extract (Psidium Guajava L ) as antioxidant that could protect liver cell membranes damage due to exposure of cigarette smoke. This research proved that red guava extract (Psidium Guajava L) could improve cell membrane that was seen through how big the serosis in liver cell in white mice after being induced by cigarette smoke. Result of conducted research by Anam(2008) proved that the poisoning level of farmers by pesticides in Batu Mediri sub- village, Karang Pule sub-district showed that from 11 farmers who did not use personal protective equipment, the 10 farmers of them (90%) suffered from mild poisoning and 1 farmer suffered from moderate poisoning. There were 13 farmers who were landowners and 112 farm laborers in Kembang Kuning Hamlet Sub-Village, Gerimax Village, Narmada Sub-District, West Lombok District, Indonesia used pesticides in organophosphate and carbamate category and type of Parathion and Selvin by spraying. Until recently, it has no research which explains that ginger rhizome can repair liver cell damage due to exposure of pesticide smoke. Moreover, according to the background above, we were interested to conduct a research with title: "Hipoprotektor ekstrak jahe sebagai antioksidan pada mencit yang terpapar pestisida" (Hypoprotector of ginger extract as an antioxidant on exposed mice by pesticides). This research was as an alternative of solution for the farmers who were pesticide poisoning by using ginger rhizome as the antioxidant because from preliminary research, it stated that ginger extract (zinger officinle rosue) as antioxidant had succeeded to improve the activity of cholinesterase enzymes and reduce SGOT (serum glutamic oxaloacetic transaminase) levels in experimental animals. From the previous research, Lilis S 2005, proved the biggest content in ginger rhizome was gingerol 6,1% and shogaol 4,3%, which were as active substances as the antioxidant. Furthermore, [this research aimed at analyzing the influence of providing ginger extract \(Zingiber officinale Roscoe\) as an antioxidant against the protection of liver and pulmonary damage due to exposure of pesticides](#) in 0,005 gr/l on white mice. RESEARCH METHOD [This research was quasi-experimental research by using simple experimental design \(Post Test Only Control Group Design\)](#). One [group in this research was](#) given a treatment and another group was not given any treatment. For the sample collection, the researchers used simpler and omsampling method. The [sample in this research was male white mice](#) which were 3-4 weeks old and their weight were about 20-25gr. The minimum sample that was used was 7 mice for each group, thus, total sample in this research was 28

mice. Schematically, design of this research could be illustrated as below:

Notes: R: randomization K0: pretest group ( without any treatment) K1 : treatment group 1( being given exposure of pesticide smoke in 0,001gr/l) K2: treatment group 2 (being given pesticide smoke and given treatment of ginger extract in 0,0001 gr/l) K3: treatment group 3 (being given pesticide smoke and given treatment of ginger rhizome extract in dose of 0,005 g/l) Research Procedures Preparation for Experimental Animal: Experimental animals (white mice) were adapted for 7 days by feeding, drinking, and resting naturally in a cage (being bred in Biochemistry Laboratory of Airlangga University in Surabaya, East Java Province, Indonesia). Besides, the mice were grouped randomly to be 4 treatment groups. Providing Pesticides in Experimental Animal: The exposure of cigarette smoke was for 1 month. In a day, the experimental mice were exposed by pesticides in 0,001 gr/l. The exposure was conducted by using nebulizer (usually was used for baby), then, it was connected to spray tool which was sprayed into the Indonesian traditional drum (kendang). Providing ginger extract: (Zingiber officinale Roscoe) was given through sonde, into white mice's mouth every day with the dose in 0,001 and 0,005 gr/l. Taking the pulmonary and liver organ from experimental animal (white mice) was conducted by surgery firstly. Some organ that will be made for microscope observation object (preparat) was saved firstly in PFA solution (Paraformaldehyd 4%). Producing microscope observation object (preparat) used HE(Hematoksilin Eosin) staining method. The observation was focused on smooth muscle tissue (conducted in Anatomy laboratory of Airlangga University Surabaya). [International Journal of Current Research, Vol. 12, Issue, 01, pp. 9281-9284, January, 2020](#)

Observation of Histopathology of Liver: Observation was conducted by looking at the microscope observation object (preparat) of histopathology of liver between treatment group and control group through magnification of 400 times. Then, it was counted for average of weight score for histopathological change. Meanwhile, the criteria of assessment of cell histopathology degrees were: Observation of Histopathology of Liver: The observation was seen from the microscope observation object (preparat) from staining result in magnification of 100x and the target that was read was the percentage of widening of alveolar lumen in broad field of view and presence of infiltration of inflammation cell.

RESEARCH RESULT

Histopathological damage of mice's (Mus musculus) Liver Tissue due to the exposure of pesticide in 0,0001 gr/l: One of the purposes in this research was in order to investigate cell damage of liver tissue as the influence of pesticide exposure in 0,001 gr/l. An observation was conducted in Control Group (K0) and Treatment Group (group that was exposed by pesticides (K1)). However, there was a significant difference from both of them. In K0 group, the average of weight score was 73% and it was occurred histopathic changes in the liver from four fields of view. Meanwhile, the average in K1 group resulted 58%. According to the observation result in K0 group, hepatocytes were seemed normal which were showed by the composition of liver cells and they were consisted of hexagonal units, which were known as lobules hepaticus (liver). The center of each lobule was a central vein, which was radially surrounded by a plate of liver cells and the shape of the cells were round and oval.

Rate of Change Value

Normal	Damage of mild hepatocyte cells ( $\leq 25\%$ )	Damage of moderate hepatocyte cells (25-50%)	Damage of chronic hepatocyte cells ( $\geq 50\%$ )
0	1	2	3

Rate of Change Value

Normal	Occurred damage cells/ infiltration of inflammation cell (mild) $\leq 25\%$ from broad field of view	Occurred damage cells/ infiltration of inflammation cell (moderate) 25-50% from broad field of view	Occurred damage cells/ infiltration of inflammation cell (chronic) $\geq 50\%$ from broad field of view
0	1	2	3

The damage of liver cell in K1 group (treatment 1) that was exposed by pesticides in 0,001 gr/l showed the liver cell degenerated and underwent necrosis in around 25 – 50%. The damage of liver cell by pesticides was likely occurred as the impact of several factors, dose, exposure time, and its types. Commonly,

the liver cell had high regeneration ability, tissue loss due to toxic substances that triggered the mechanism of cell division and it continued until the improvement of tissue mass was achieved. Furthermore, necrosis process was begun by weakening of blood vessel wall, inflammatory response, and the formation of plaque which was foam cells and the blood clots was formed. Foam cell was a layer that contained macrophages which was laden with lipid and smooth muscle cells whose cytoplasm was raised by lipids. This foam cell was a component in forming lipid zones and smooth muscle cells that enlarged by lipids. Fatty streak was consisted of fat-filled foam cells which were lesions that did not increase significantly, thus, it did not cause impaired blood flow. Furthermore, the correlation between fatty streak and atherosclerotic plaques was unclear although these patches could develop into plaques. Improvement of liver tissue cell as the impact of giving ginger extract (*Zingiber officinale* Roscoe) in dose of 0,0001 gr/l against cell damage in liver organ due to pesticide exposure in dose of 0,0005 gr/l: In K2 group (treatment 2 (it was exposed by pesticide and was given [ginger extract \(\*Zingiber officinale\* Roscoe\)\) as antioxidant against the improvement of tissue cell in histopathology of broncus that underwent cell damage due to exposure of cigarette smoke in white mice. The description of liver cell in this K2 group was seemed to degenerate cell. Degeneration change was a change whose process was reversible, which meant that it could return as what it was in the beginning. Thus, cell would return to be healthy as before being exposed by pesticides. Theoretically, the process of liver cell damage was begun from degeneration process with criteria of cell swelling. The criterion was observed in the treatment \[of providing ginger extract \\(\\*Zingiber officinale\\* Roscoe\\)\]\(#\) that was \[as antioxidant\]\(#\). A treatment \[of providing gingerol, shogaol, dheartil heptanoite as high antioxidant\]\(#\) caused extracellular fluid entered into the cytosol in large quantities. \(Annisa, Agata 2016\). In K2 group that was given ginger extract \(\*Zingiber officinale\* Roscoe\) as antioxidant was obtained data that damage of histopathological structure decreased more than in K1 group \(without any \[providing ginger extract \\(\\*Zingiber officinale\\* Roscoe\\)\]\(#\)\). Shogaol that was \[as an antioxidant\]\(#\) had role in improving liver cell damage due to pesticide exposure with recovery effect and the level of damage was mild \(0- 25%\). Improvement of tissue cell in liver due to providing ginger extract\(\*Zingiber officinale\* Roscoe\) in dose of 0,005gr/l against cell damage in liver due to pesticide exposure in dose of 0,0001 gr/l: In K3 group was different with K1 group which had been exposed by pesticide in 0,0001 gr/l for 30 days and it was injected with ginger extract as the antioxidant. Description of histopathological liver in K3 group was seen that the damage of histopathological structure decreased more rather than in K1 group \(treatment only with pesticide\). In this group, there was hepatosis that underwent microvesicular steatosis \(fatty degeneration\). Besides, in liver cell, a central vein underwent blood containment that was seen as lumen of central vein and it was covered by a red mass which was the blood that blocked these vessels. Besides, there were also bleeding of veriver and inflammation cells around the liver cell. In this K3 group, there were central vein, hepatocyte, and lobius cells which were almost in normal condition. Although the decrease of damage had not reached the illustration of normal histopathology, result of this research had undergone cell improvement and this was possibly caused by lack of time in providing ginger extract \(\*Zingiber officinale\* Roscoe\) or a mistake in providing the dose. Compound in category of Shogaol an gingerol also could induce necrosis and stopped the cell cycle through the mechanism of inhibition of enzyme topoisomerase. Shogaol an gingerol also improved excretion of the glutathione S -transferase enzyme which could detoxify carcinogens so that the body was eliminated quickly. Conclusion ? There was damage of the structure of liver tissue \(moderate hepatocyte damage\) and pulmonary \(scar Umi Rahayu et al. \[Potency of ginger extract \\(\\*zingiber officinale\\* roscoe\\) as antioxidant and hepatoprotector for liver and pulmonary tissue in exposed mice \\(mus\]\(#\)](#)

[musculus\) by organophosphate pesticides](#) tissue) from the white mice due to pesticide exposure in Guyton, A.C. & John E.H, 2007. Text Book of Medical 0,1. 10 -6 gr/lit Physiology, 11th edition. Elsevier Saunders, 1600 John F. ? The improvement of damaged liver cell was through Kennedy Blvd., Suite 1800. Philadelphia, Pennsylvania, pp. cell degeneration and the cell damage was in mild – 1063-1072, 1129-1132, and 1339-1347. moderate. In pulmonary organ, scarring had not Hana. [2010. Lama Pajanan organofosfat terhadap penurunan](#) occurred anymore. [aktivitas enzim kolinesterase dalam darah petani sayuran](#). ? The improvement of liver tissue cells due to providing [MAKARA Kesehatan Vol 14](#). no2. [Desember 2010](#). ginger extract (Zingiber officinale Roscoe) in dose of Harris, L.E. 1970. Nutrition Research Tehnique for Domestic 0,0005 gr/l against cell damage in liver due topesticides and Wild Animals. Departemen Animal Nutrition Utah exposure in 0,0001 gr/lit showed cell degeneration and State University Logan, Utah. hepatotic damage, such as damage with hepatocyte Hayati, A., Yunaida, B., Pidada, R., Darmanto, W., dan score in mild, moderate and chronic. Winarni, D., 2004. Efek 2 Methoxyethanol Terhadap Struktur Histologi Mencit (Mus musculus). Berkala Recommendation Penelitian Hayati, 10(1): 7-12 Hukkanen, J., Jacob III, P. & Benowitz, N.L. 2005. The results of this study can be used to repair lung organ cells [Metabolism and Disposition Kinetics of Nicotine](#). [The for](#) people who use pesticides in agricultural practices or pest American Society for Pharmacology and Experimental eradication. Need to do research on the administration of Therapeutics. Vol. 57, No. 1. ginger extract in mice exposed to pesticides with different McPhee, S.J & Pignone, M. 2007. Disease Prevention and types of ginger, dosage and time. Health Promotion. In S.J. McPhee, M.A, Papadakis, & L.M. Tierney Jr (Eds), Current Medical Diagnosis and REFERENCES [Treatment, 47th Edition, \(p. 1-16\)](#). New York: McGraw- Hill. Achmad, F., 2010. Kadar SGPT dan gambaran Histopatologis Ni Putu, M2015. Uji aktivitas antibakteri ekstrak Etanol Hepar akibat paparan alletrin, Skripsi Universitas Negeri rimpang jahe merah terhadap bakteri Klebsiella pneumonie Malang. isolate sputum penderita bronchitis secara in vivo. Jurnal Ali, M, Yuyun, Y.2004. Pengaruh Paparan asap rokok Kretek Ilmiah Farmasi UNSRAT Vol4 no.3. Agustus 2015 Sadikin terhadap Peroksidasi Lemak dan System proteksi 2003, Antioksidan, Radikal bebas dan Superoksida Dismutase Hepar Tikus Wistar . Jurnal penuaan. www.chem-is.try.com Kedokteran YARSI Sofia, 2006. Antioksidan dan Radikal .www.chem-is.try.com Annisa, Agata 2016. Respon histopatologis Hepar mencit (mus Sudarsono,P .2002. Tumbuhan Obat Ondonesia .Yogyakarta musculus) yang diinduksi benzopiren terhadap pemberian Uikey AK, VK Hazare SM, and Vaidhya. 2003. Estimation of taurin dan ekstrct daun sirsat (annona Muricata), Jurnal serum antioxidant enzymes SOD dan GPx in oral Nature Indoneia 16(2), April 2016: 54-62. submucus fibrosis : A Biochemichal Study. Journal of Oral Departemen Kesehatan RI. 2004. Kawasan TanpaRokok.Pusat &Maxillofacial Pathology:Vol 7 Issue 2 p:44-45. Promkes Depkes RI. Jakarta. Utami ,Aprihatin, dkk. 2012. Variasi metode DNA daun Evans,W. J.2000.VitaminE, vitaminC, and exercise. Am J Clin temulawak (curcuma xanthori). PROSEDING Seminar Nutr. Nasional Kimia Unesa ISBN 978-979-028 550-7 Fidrianny, I.,I. Supradja., dan A. Soemardji. 2004. Wang HO, Reilly EJ, Weisskopf MG, Logroscino G, Analisis nikotin dalam asap dan filter rokok. Acta McCullough, M.L Thun M J. 2011. Smoking and risk of Pharmaceutica Indonesia; 29(3):100 amyotrophic lateral sclerosis: A pooled analysis of 5 Rokhmalia, F. 2016. Peningkatan Aktivitas Enzim SOD Serum prospective cohorts.. Arch Neurol.,68(2): 207-215. Dan Keluhan Kesehatan Terhadap Paparan Asap Widodo, E. Pajanan Asap Rokok Kretek Pada Tikus Putih Pembakaran Kayu Pada Pekerja”. Suara Forikes, April Sebagai Model Untuk Manusia. Bogor: Sekolah 2016:98-102. Pascasarjana Intitut Pertanian Bogor; 2006. 2-3. Freisleben. H.J. 2001. Free radicals and ROS (Reactive Winarsi, H.2007. Antioksidan Alamidan Radikal Bebas Oxygen Species) in biological systems. Dalam : Kursus

Potensidan Aplikasi dalam Kesehatan.Yogyakarta. penyegar dan pelatihan  
2001 radikal bebas dan antioksidan Kanisius. dalam kesehatan dasar,  
aplikasi, dan pemanfaatan bahan Yuli,H. 2016. Pemanfaatan ekstrak air  
rimpang jahe merah alam. Jakarta: Bagian Biokimia FKUI. hal. 1-21  
Ganong, (Zingiber offcinnale) pada Biosintesis sederhana nano W.F. 2002.  
Buku ajar fisiologi kedokteran. Edisi 20. partikel perak. Jakarta Gildea, TR,  
2009. McCarthy K, Pothier C, Kavuru MS, Lauer MS. Association of an  
abnormal exercise heart rate recovery with pulmonary function  
abnormalities. Chest 2004; 125:1286-91 \*\*\*\*\* 9282 9283 9284